arXiv:1201.4669v2 [math.CO] 27 Apr 2012

ON MAXIMAL CHAINS IN THE NON-CROSSING PARTITION LATTICE

RON M. ADIN AND YUVAL ROICHMAN

Abstract. A weak order on the set of maximal chains of the non-crossing partition lattice is introduced and studied. A zero-Hecke algebra action is used to compute the radius of the graph on these chains in which two chains are adjacent if they differ in exactly one element.

Contents

1. Introduction 1
2. Basic concepts 2
3. A map from F_n to S_{n-1} 4
4. Properties of the weak order on F_n 7
5. Zero-Hecke algebra action 8
6. Number of maximal elements 12
7. Right and left inversions 14
8. Right inversions and q-Catalan numbers 17
9. Radius and diameter 21
References 23

1. Introduction

Consider the graph $G_T(n)$ with vertex set consisting of all maximal chains in $NC(n)$, the non-crossing partition lattice of type A, and where two chains are adjacent if they differ in exactly one element. This graph may be identified with the graph of all reduced words of a given Coxeter element (long cycle) in the symmetric group, where the alphabet consists of all reflections (transpositions) and two words are adjacent if they agree in all but two adjacent letters, which are tr in one word and $r^t t$ or rt^r in the other. Here $g^h := h^{-1}gh$. This graph is known to be connected – see, e.g., [3, Prop. 1.6.1]; we are interested in calculating

---

Date: April 27, '12.

Key words and phrases. Coxeter group, symmetric group, zero-Hecke algebra, reduced word, weak order, radius, Catalan number, Hurwitz action.

Reserach of both authors was partially supported by an internal grant from Bar-Ilan University.

its radius. This is motivated by [2] and [14], where the analogous question for simple reflections was studied, and by [11] which evaluated the radius of a related graph on labeled trees. Recall that, by a classical result of Hurwitz [12, 7, 17], the number of maximal chains in the non-crossing partition lattice of type A is equal to the number of labeled trees on n vertices.

Our approach is to consider a zero-Hecke algebra action on the set of maximal chains. This allow us to define a well-behaved natural weak order on this set. Each maximal interval is isomorphic to the weak order on the symmetric group. The number of maximal elements, refined by a generalized inversion number, is the Carlitz-Riordan q-Catalan number. The resulting undirected Hasse diagram spans the graph $G_T(n)$ of maximal chains, implying an evaluation of the radius and an approximation of the diameter up to a factor 3/2.

2. Basic concepts

The non-crossing partition lattice $NC(n)$, first introduced by Kreweras [13], may be defined as follows.

Let T be the set of all reflections (transpositions) in the symmetric group S_n, and let $\ell(\cdot)$ be the corresponding length function: $\ell(\pi)$ is the minimal number of factors in an expression of π as a product of reflections. Let c be a Coxeter element in this group (i.e., a cycle of length n); for concreteness, take $c = (1, 2, \ldots, n)$. Then $NC(n)$ is the set

$$\{\pi \in S_n \,:\, \ell(\pi) + \ell(\pi^{-1}c) = \ell(c)\}$$

ordered by

$$\pi \le \sigma \iff \ell(\pi) + \ell(\pi^{-1}\sigma) = \ell(\sigma).$$

Definition 2.1. Let F_n be the set of all maximal chains in the non-crossing partition lattice $NC(n)$.

Clearly, an element of F_n corresponds to a factorization of c into a minimal number ($\ell(c) = n - 1$) of transpositions. It can thus be written in the form $(t_1, \ldots, t_{n-1})$, where $t_i \in T$ $(1 \le i \le n-1)$ and $t_1 \cdots t_{n-1} = c$.

Definition 2.2. The *Hurwitz graph* $G_T(n)$ is the (undirected) graph with vertex set F_n, where two chains are adjacent if they differ in exactly one element.

The Hurwitz graph $G_T(4)$ is drawn in Figure 2.1; note that $|F_4| = 16$, and the midpoint of of the figure is not a vertex.

Definition 2.3. For each $1 \le i \le n-2$ and $v = (t_1, \ldots, t_{n-1}) \in F_n$ let

$$R_i(v) := (t_1, \ldots, t_{i-1}, t_{i+1}^{t_i}, t_i, t_{i+2}, \ldots, t_{n-1})$$

and

$$L_i(v) := (t_1, \ldots, t_{i-1}, t_{i+1}, t_i^{t_{i+1}}, t_{i+2}, \ldots, t_{n-1}),$$

where $g^h := h^{-1}gh$.



FIGURE 2.1. The Hurwitz graph $G_T(4)$

R_i and L_i $(1 \leq i \leq n-2)$ are operators on F_n, where $R_i(v)$ slides the i-th factor to the right while $L_i(v)$ slides the $(i+1)$-st factor to the left. Clearly $R_i = L_i^{-1}$ and, for every $v \in F_n$, either $R_i^2(v) = v$ (if t_i and t_{i+1} commute) or $R_i^3(v) = v$ (otherwise). Furthermore, the following is true.

Proposition 2.4. [12] *For every* $1 \leq i \leq n-2$,

$$R_i R_{i+1} R_i = R_{i+1} R_i R_{i+1}.$$

This clearly defines a braid group action on F_n, a distinguished example of the Hurwitz action on ordered tuples of group elements.

Observation 2.5. For $u, v \in F_n$, $\{u, v\}$ is an edge of $G_T(n)$ if and only if there exists an $1 \leq i \leq n-2$ such that either $v = R_i(u)$ or $v = L_i(u)$.

Remark 2.6. The classical Tits graph of a Coxeter group is the graph whose vertices are all the reduced words of a longest element of the group, with respect to the set of *simple* reflections S; two words are adjacent if one is obtained from the other by a braid relation. In other words, vertices correspond to maximal chains in the weak order on the group; two chains are adjacent if they differ in a “minimal cycle”. Replacing the set S of simple reflections by the set T of all reflections, one obtains the Hurwitz graph. The diameters of the Tits graphs of the Coxeter groups of types A and B were evaluated in [14]. Bessis proved that the Hurwitz action on F_n is transitive [3, Prop. 1.6.1], thus that the Hurwitz graph $G_T(n)$ is connected. Its radius and bounds on its diameter will be given in Section 9.

Consider the following definition of a partial order on F_n. A similar definition applies to any finite connected graph with a distinguished vertex, endowing the set of vertices (in our case, F_n) with the structure of a graded poset.

Definition 2.7. For $u, v \in F_n$, denote $u \leq v$ if u belongs to a geodesic in $G_T(n)$ from $e := ((1,2),(2,3),\ldots,(n-1,n))$ to v. The resulting order will be called *the weak order on* F_n, and the resulting poset will be denoted by $\mathrm{Weak}(F_n)$.

The Hasse diagram of $\mathrm{Weak}(F_4)$ is drawn in Figure 2.2. An edge is labeled (colored) i if it corresponds to an action of R_i (or of L_i, in the opposite direction), for $1 \leq i \leq 2$.



FIGURE 2.2. The Hasse diagram of $\mathrm{Weak}(F_4)$

Remark 2.8. By Proposition 2.4, $G_T(n)$ is a Schreier graph of the braid group of type A, with respect to a suitable subgroup. Thus, by definition, the weak order on the braid group, restricted to the shortest coset representatives, is isomorphic to $\mathrm{Weak}(F_n)$.

Properties of the poset $\mathrm{Weak}(F_n)$ are studied in this paper.

3. A MAP FROM F_n TO S_{n-1}

Let F_n be as in Section 1. Define a map $\phi : F_n \to S_{n-1}$ as follows.

Definition 3.1. For $w = (t_1, \ldots, t_{n-1}) \in F_n$ define the partial products

$$\sigma_j := t_j t_{j+1} \cdots t_{n-1} \qquad (1 \leq j \leq n-1)$$

and the empty product

$$\sigma_n := id.$$

By definition, $\sigma_j = t_j\sigma_{j+1}$ for every $1 \le j \le n-1$. Thus $\sigma_j(i) \ne \sigma_{j+1}(i)$ for exactly two values of $1 \le i \le n$, and $\sigma_j(i) > \sigma_{j+1}(i)$ for exactly one such value. Ignoring $i = n$, the set

$$A_j := \{1 \le i \le n-1 \,|\, \sigma_j(i) > \sigma_{j+1}(i)\}$$

thus satisfies

$$|A_j| \le 1 \qquad (1 \le j \le n-1). \tag{1}$$

On the other hand, $\sigma_1(i) = c(i) = i+1$ while $\sigma_n(i) = i$, so that $i \in \bigcup_{j=1}^{n-1} A_j$ for every $1 \le i \le n-1$. It follows that

$$\left|\bigcup_{j=1}^{n-1} A_j\right| \ge n-1. \tag{2}$$

Combining (1) with (2) one concludes that

$$\left|\bigcup_{j=1}^{n-1} A_j\right| = n-1$$

and

$$|A_j| = 1 \qquad (\forall j).$$

It follows that the map π_w defined by

$$\pi_w(j) := i \text{ if } A_j = \{i\}$$

is a permutation in S_{n-1}.

Definition 3.2. Define $\phi : F_n \to S_{n-1}$ by

$$\phi(w) := \pi_w \qquad (\forall w \in F_n).$$

Observation 3.3. For every $w \in F_n$ and $1 \le j \le n-1$, if $t_j = (a, b)$ with $a < b$ then

$$\phi(w)(j) = \sigma_{j+1}^{-1}(a).$$

Example 3.4. Let $w = ((2,5),(1,5),(2,4),(2,3)) \in F_5$. Then $\sigma_5 = id$, $\sigma_4 = (2,3)$, $\sigma_3 = (2,3,4)$, $\sigma_2 = (2,3,4)(1,5)$ and $\sigma_1 = (1,2,3,4,5) = c$, where permutations are written in cycle notation. Thus $\phi(w)(1) = 4 = \sigma_2^{-1}(2)$, $\phi(w)(2) = 1 = \sigma_3^{-1}(1)$, $\phi(w)(3) = 3 = \sigma_4^{-1}(2)$ and $\phi(w)(4) = 2 = \sigma_5^{-1}(2)$.

For a sequence $v = ((a_1, b_1), \ldots, (a_\ell, b_\ell))$ of ℓ transpositions in S_n, let $G(v)$ be the geometric graph whose vertices are $\{1, \ldots, n\}$ drawn on a circle and whose edges are $\{(a_i, b_i) : \ 1 \le i \le \ell\}$. The following characterization of the words in F_n, due to Goulden and Yong, is very useful.

Proposition 3.5. [10, Theorem 2.2]
For $t_1, \ldots, t_{n-1} \in T(S_n)$, the word $v = (t_1, \ldots, t_{n-1})$ belongs to F_n if and only if the following three conditions hold:

(i) *$G(v)$ is a tree.*

(ii) *$G(v)$ is non-crossing; namely, two edges may intersect only in a common vertex.*

(iii) (cyclic order on neighbors) *For every $1 \le a \le n$ and $1 \le i < j \le n-1$, if $t_i = (a,c)$ and $t_j = (a,b)$ then $b <_a c$. Here $<_a$ is the linear order $a <_a a+1 <_a \cdots <_a n <_a 1 <_a \cdots <_a a-1$.*

Proposition 3.5(iii) may be stated in the following equivalent form.

Corollary 3.6. *For $v = (t_1, \ldots, t_{n-1}) \in F_n$ and $i < j$, if t_i and t_j do not commute then (t_i, t_j) is either $((a,c),(a,b))$, $((b,c),(a,c))$ or $((a,b),(b,c))$, for some $a < b < c$.*

For integers $a < b$ denote: $[a,b) := \{a, a+1, \ldots, b-1\}$. Proposition 3.5 together with Observation 3.3 imply the following two lemmas.

Lemma 3.7. *For $v = (t_1, \ldots, t_{n-1}) \in F_n$, if $t_i = (a,b)$ with $a < b$ then $\phi(v)(i) \in [a,b)$.*

Lemma 3.8. *For $v = (t_1, \ldots, t_{n-1}) \in F_n$ and $i \ne j$, if $t_i = (a,d)$ and $t_j = (b,c)$ with $a < b < c < d$ then*

(1) *$\phi(v)(i) \notin [b,c)$.*

(2) *Deleting the edge (a,d) from the tree $G(v)$ leaves two connected components: T_a (containing a) and T_d (containing d). Then $\phi(v)(i) \in [c,d)$ if and only if $T_a \cap [c,d) \ne \emptyset$.*

Lemma 3.9. (characterization of inversions in $\phi(v) \in S_{n-1}$)
For $v = (t_1, \ldots, t_{n-1}) \in F_n$ and $i < j$, $\phi(v)(i) > \phi(v)(j)$ if and only if one of the following holds:

(1) *$\exists a < b < c$ such that either:*
    (i) *$t_i = (a,c)$ and $t_j = (a,b)$; or:*
    (ii) *$t_i = (b,c)$ and $t_j = (a,c)$.*

(2) *$\exists a < b < c < d$ such that $t_i = (c,d)$ and $t_j = (a,b)$.*

(3) *$\exists a < b < c < d$ such that either:*
    (i) *$t_i = (a,d)$, $t_j = (b,c)$ and the graph with edges $\{t_{i+1}, \ldots, t_{n-1}\}$ contains a path from a to some $x \in [c,d)$; or*
    (ii) *$t_j = (b,c)$, $t_j = (a,d)$ and the graph with edges $\{t_{i+1}, \ldots, t_{n-1}\}$ does not contain a path from a to any $x \in [c,d)$.*

Proof. (2) follows from Lemma 3.7. (1) and (3) follow from Lemma 3.7 together with both parts of Lemma 3.8.

□

Recall the operators R_j and L_j from Section 2. By definition, if we represent a word $w \in F_n$ by the sequence of partial products $(\sigma_1, \ldots, \sigma_n)$, then for all $1 \le j \le n-2$

$$R_j(\ldots, \sigma_j, \sigma_{j+1}, \sigma_{j+2}, \ldots) = (\ldots, \sigma_j, t_j t_{j+1} \sigma_{j+1}, \sigma_{j+2}, \ldots), \tag{3}$$

and

$$L_j(\ldots,\sigma_j,\sigma_{j+1},\sigma_{j+2},\ldots) = (\ldots,\sigma_j,t_{j+1}t_j\sigma_{j+1},\sigma_{j+2},\ldots). \tag{4}$$

Lemma 3.10. *For every* $w \in F_n$ *and* $1 \le j \le n-2$,

$$\phi(R_j(w)) = \begin{cases} \phi(w), & \text{if } \exists\, a<b<c \ \text{ s.t. } t_j=(a,c) \text{ and } t_{j+1}=(a,b); \\ \phi(w)s_j, & \text{otherwise} \end{cases} \tag{5}$$

and

$$\phi(L_j(w)) = \begin{cases} \phi(w), & \text{if } \exists\, a<b<c \ \text{ s.t. } t_j=(b,c) \text{ and } t_{j+1}=(a,c); \\ \phi(w)s_j, & \text{otherwise.} \end{cases} \tag{6}$$

Proof. We prove (5). The proof of (6) is similar.

By (3), R_j affects, among the partial products of w, only σ_{j+1}. It follows, by Observation 3.3, that $\phi(R_j(w))(i) = \phi(w)(i)$ for all $i \notin \{j, j+1\}$. Thus either $\phi(R_j(w)) = \phi(w)$ or $\phi(R_j(w)) = \phi(w)s_j$. By Observation 3.3 (for $j+1$), $\phi(R_j(w)) = \phi(w)$ if and only if $t_{j+1} = (a,b)$ and $t_j = (a,c)$ for some a, b and c such that $a < b$ and $a < c$. These t_j and t_{j+1} do not commute, and by Corollary 3.6 necessarily $b < c$ as well.

□

4. Properties of the weak order on F_n

Recall the graded poset Weak(F_n) from Definition 2.7. Note that, by definition,

Fact 4.1. The word $e = ((1,2),(2,3),\ldots,(n-1,n))$ is the unique minimal element in Weak(F_n).

Lemma 4.2.

$$\phi^{-1}(id) = \{e\}.$$

Proof. Clearly $\phi(e) = id$, since for $e = ((1,2),\ldots,(n-1,n))$ one has $t_j = (j,j+1)$, $\sigma_j = t_j \cdots t_{n-1} = (j,\ldots,n)$ and thus $\phi(e)(j) = \sigma_{j+1}^{-1}(j) = j$ by Observation 3.3.

Conversely, assume that $w = (t_1,\ldots,t_{n-1}) \in F_n$ has $\phi(w) = id$. Then $\phi(w)(n-1) = n-1$ implies that $\sigma_{n-1}(n-1) > \sigma_n(n-1) = n-1$, so that $\sigma_{n-1}(n-1) = n$ and $t_{n-1} = \sigma_{n-1} = (n-1,n)$. It follows that $t_1 \cdots t_{n-2} = (1,\ldots,n)(n-1,n) = (1,\ldots,n-1)$, so that necessarily $t_1,\ldots,t_{n-2} \in S_{n-1}$ and $w' := (t_1,\ldots,t_{n-2})$ belongs to F_{n-1}, with $\phi(w') = id \in S_{n-2}$. Induction on n completes the proof.

□

For any $w \in F_n$, let rank(w) be its rank in the poset Weak(F_n). For $\pi \in S_{n-1}$, let inv(π) be its inversion number.

Lemma 4.3. *For any* $w \in F_n$,

$$\operatorname{rank}(w) = \operatorname{inv}(\phi(w)).$$

Proof. By induction on $\mathrm{inv}(\phi(w))$.

If $\mathrm{inv}(\phi(w)) = 0$ then $\phi(w) = id$. By Lemma 4.2 $w = e$, so that indeed $\mathrm{rank}(w) = 0 = \mathrm{inv}(\phi(w))$.

For the induction step let $\mathrm{inv}(\phi(w)) = \ell > 0$, and assume that the claim holds for $u \in F_n$ whenever $\mathrm{inv}(\phi(u)) < \ell$. Let $\pi := \phi(w) \in S_{n-1}$. Since $\mathrm{inv}(\pi) > 0$ there exists $1 \le j \le n-2$ such that $\mathrm{inv}(\pi s_j) < \mathrm{inv}(\pi)$. By Lemma 3.10, $\phi(R_j(w)) = \pi s_j$ unless $\exists\, a < b < c$ such that $t_j = (a,c)$ and $t_{j+1} = (a,b)$ (in w); whereas $\phi(L_j(w)) = \pi s_j$ unless $\exists\, a < b < c$ such that $t_j = (b,c)$ and $t_{j+1} = (a,c)$. These two exceptions cannot hold simultaneously, and thus $\phi(u) = \pi s_j$ for either $u = R_j(w)$ or $u = L_j(w)$. For such u, $\mathrm{inv}(\phi(u)) = \mathrm{inv}(\pi s_j) = \ell - 1$, so by the induction hypothesis $\mathrm{rank}(u) = \mathrm{inv}(\phi(u))$. Since u and w are connected by an arc in the graph $G_T(n)$, it follows that $\mathrm{rank}(w) \le \mathrm{rank}(u) + 1 = \ell$. On the other hand, in a geodesic from e to w in $G_T(n)$ the value of $\mathrm{inv}(\phi(\cdot))$ changes by 0 or 1 on each arc (again by Lemma 3.10), so that $\mathrm{rank}(w) \ge \mathrm{inv}(\phi(w)) = \ell$. Thus $\mathrm{rank}(w) = \ell = \mathrm{inv}(\phi(w))$.

□

Corollary 4.4. *The (undirected) Hasse diagram of* $\mathrm{Weak}(F_n)$ *is obtained from the graph* $G_T(n)$ *by deleting all edges connecting a vertex* $(\ldots,(a,c),(a,b),\ldots)$ *with a vertex* $(\ldots,(b,c),(a,c),\ldots)$ *for some* $a < b < c$.

Proof. By Lemma 3.10 and Lemma 4.3. □

Lemma 4.5. *For every* $v \in F_n$ *and* $1 \le j \le n-2$*, if* $R_j(v) < v$ *and* $L_j(v) < v$ *in* $\mathrm{Weak}(F_n)$ *then* $R_j(v) = L_j(v)$.

Proof. Assume that $R_j(v) < v$ and $L_j(v) < v$. We can assume that t_j and t_{j+1} do not commute (otherwise, by definition, $R_j(v) = L_j(v)$). Then, by Corollary 3.6, $(t_j, t_{j+1}) \in \{((a,c),(a,b)),\ ((b,c),(a,c)),\ ((a,b),(b,c))\}$.

By Lemma 4.3, $R_j(v) < v$ implies that $\mathrm{inv}(\phi(R_j(v))) < \mathrm{inv}(\phi(v))$, which is equivalent, by Lemma 3.10, to $\phi(R_j(v))(j) < \phi(R_j(v))(j+1)$. By Lemma 3.9(1), this happens if and only if the j-th factor in $R_j(v)$ is (a,b) and the $(j+1)$-st factor is (b,c). By similar arguments, $L_j(v) < v$ implies that the j-th factor in $L_j(v)$ is (a,b) and the $(j+1)$-st factor is (b,c). Thus $R_j(v) = L_j(v)$.

□

Remark 4.6. Lemma 4.5 may be stated in the following two equivalent forms.

- For every $v \in F_n$ and $1 \le j \le n-2$, $R_j(v) > v$ if and only if $L_j(v) > v$.
- The orbit in F_n of each R_j (or L_j) consists of either a pair $\{v_1, v_2\}$ with $v_1 < v_2$ or a triple $\{v_1, v_2, v_3\}$ with $v_1 < v_2$ and $v_1 < v_3$.

5. Zero-Hecke algebra action

The *zero-Hecke algebra* $\mathcal{H}_n(0)$ is an algebra over $\mathbb{Q}$ generated by $\{T_i : 1 \le i \le n-1\}$ with defining relations

$$T_i^2 = T_i \qquad (1 \le i \le n-1), \tag{7}$$

$$T_iT_j = T_jT_i \qquad (|i-j|>1) \tag{8}$$

and

$$T_iT_{i+1}T_i = T_{i+1}T_iT_{i+1} \qquad (1 \le i \le n-2). \tag{9}$$

$\mathcal{H}_n(0)$ can be viewed as the specialization $q = 0$ of the Iwahori-Hecke algebra $\mathcal{H}_n(q)$, deforming the group algebra of S_n, or as the semigroup algebra of a suitable monoid. Note that our notation is slightly non-standard, since equation (7) is usually stated as $T_i^2 = -T_i$, which amounts to replacing T_i by $-T_i$ (for all i).

A faithful action of the zero-Hecke algebra $\mathcal{H}_{n-1}(0)$ on F_n is introduced in this section. This action will be applied to show that every maximal interval in $\text{Weak}(F_n)$ is isomorphic to the weak order on the symmetric group S_{n-1}.

5.1. **Down operators.** Recall the descent set of a permutation $\pi \in S_{n-1}$, $\text{Des}(\pi) := \{i : \ \pi(i) > \pi(i+1)\}$.

Definition 5.1. For $1 \le i \le n-2$ define the *i-th down operator* $D_i : F_n \longrightarrow F_n$ by

$$D_i(w) := \begin{cases} R_i(w), & \text{if } i \in \text{Des}(\phi(w)) \text{ and } R_i(w) < w; \\ L_i(w), & \text{if } i \in \text{Des}(\phi(w)) \text{ and } L_i(w) < w; \\ w, & \text{if } i \notin \text{Des}(\phi(w)). \end{cases}$$

Claim 5.2. The down operators are well defined.

Proof. By Lemma 3.10, for any $w \in F_n$ and $1 \le i \le n-2$, either $\phi(R_i(w)) = \phi(w)s_i$ or $\phi(L_i(w)) = \phi(w)s_i$ (or both). If $i \in \text{Des}(\phi(w))$ then $\text{inv}(\phi(w)s_i) < \text{inv}(\phi(w))$. Thus, by Lemma 4.3, either $\text{rank}(R_i(w)) < \text{rank}(w)$ or $\text{rank}(L_i(w)) < \text{rank}(w)$, namely: Either $R_i(w) < w$ or $L_i(w) < w$ in $\text{Weak}(F_n)$. If both hold then, by Lemma 4.5, $R_i(w) = L_i(w)$.

□

Observation 5.3. For every $1 \le i \le n-2$ and $w \in F_n$,

$$\phi(D_i(w)) = \begin{cases} \phi(w)s_i, & \text{if } i \in \text{Des}(\phi(w)); \\ \phi(w), & \text{if } i \notin \text{Des}(\phi(w)). \end{cases}$$

Lemma 5.4. *The down operators $\{D_i \ : \ 1 \le i \le n-2\}$ satisfy the defining relations of the zero-Hecke algebra $\mathcal{H}_{n-1}(0)$.*

Proof. By Definition 5.1, if $i \notin \text{Des}(\phi(w))$ then $D_i^2(w) = (w)$. If $i \in \text{Des}(\phi(w))$ then, by Observation 5.3, $i \notin \phi(D_i(w))$, hence by Definition 5.1, $D_i^2(w) = D_i(D_i(w)) = D_i(w)$. Thus $D_i^2 = D_i$ for every $1 \le i < n-1$.

If $|i-j| > 1$, then by Definition 5.1, $D_iD_j = D_jD_i$.

We have to verify that $D_iD_{i+1}D_i = D_{i+1}D_iD_{i+1}$ for every $1 \le i < n-2$.

If $i, i{+}1 \notin \text{Des}(\phi(w))$ then, by Definition 5.1, $D_iD_{i+1}D_i(w) = w = D_{i+1}D_iD_{i+1}(w)$.

If $i \in \mathrm{Des}(\phi(w))$ and $i+1 \notin \mathrm{Des}(\phi(w))$ then, by Definition 5.1, $D_{i+1}(w) = w$, thus $D_{i+1}D_iD_{i+1}(w) = D_{i+1}D_i(w)$. On the other hand, in this case

$$\phi(D_{i+1}D_i(w))(i) = \phi(D_i(w))(i) = \phi(w)(i{+}1) = \min\{\phi(w)(i), \phi(w)(i{+}1), \phi(w)(i{+}2)\}$$

$$= \min\{\phi(D_{i+1}D_i(w))(i), \phi(D_{i+1}D_i(w))(i+1), \phi(D_{i+1}D_i(w))(i+2)\}.$$

Hence $i \notin \mathrm{Des}(\phi(D_{i+1}D_i(w)))$, thus, by definition, $D_iD_{i+1}D_i(w) = D_{i+1}D_i(w)$.

For similar arguments, if $i \notin \mathrm{Des}(\phi(w))$ and $i{+}1 \in \mathrm{Des}(\phi(w))$ then $D_iD_{i+1}D_i(w) = D_iD_{i+1}(w) = D_{i+1}D_iD_{i+1}(w)$.

It remains to verify the braid relation $D_iD_{i+1}D_i = D_{i+1}D_iD_{i+1}$, when $i, i{+}1 \in \mathrm{Des}(\phi(w))$.

For $w = ((t_1, \ldots, t_{n-1}) \in F_n$ let $w_i := t_i = (a_i, b_i)$.

Obviously, if $j \notin \{i, i+1, i+2\}$ then $D_iD_{i+1}D_i(w)_j = D_{i+1}D_iD_{i+1}(w)_j = t_j$. Thus, it suffices to verify that $D_iD_{i+1}D_i(w)_j = D_{i+1}D_iD_{i+1}(w)_j$ for $j \in \{i, i+1, i+2\}$.

For $I \subseteq \{1, \ldots, n-1\}$ let $\mathrm{Supp}_I(w) := \cup_{i \in I}\{a_i, b_i\}$.

By Proposition 3.5(i), $4 \le |\,\mathrm{Supp}_{\{i,i+1,i+2\}}(w)| \le 6$.

Case 1. If $|\,\mathrm{Supp}_{\{i,i+1,i+2\}}(w)| = 6$ then t_i, t_{i+1} and t_{i+2} commute.

Then

$$D_iD_{i+1}D_i(w)_j = t_{\phi(w)^{-1}T_iT_{i+1}T_i(\phi(w))(j)}$$

with natural action of $\mathcal{H}_{n-1}(0)$ on S_{n-1} (namely, $T_i(\pi) := \pi s_i$ if $i \in \mathrm{Des}(\pi)$ and π otherwise). Similarly,

$$D_{i+1}D_iD_{i+1}(w)_j = t_{\phi(w)^{-1}T_{i+1}T_iT_{i+1}(\phi(w))(j)}.$$

Since $T_iT_{i+1}T_i = T_{i+1}T_iT_{i+1}$ we are done in this case.

Case 2. If $|\,\mathrm{Supp}_{\{i,i+1,i+2\}}(w)| = 5$ then one of the three transpositions t_i, t_{i+1} or t_{i+2} commutes with other two.

If t_i commutes with t_{i+1} and t_{i+2} then,

since $i+1 \in \mathrm{Des}(\phi(w))$, letting $\mathrm{Supp}_{\{i+1,i+2\}} = \{a, b, c\}$ with $a < b < c$, it follows from Lemma 3.9, that either $t_{i+1} = (a, c)$ and $t_{i+2} = (a, b)$ or $t_{i+1} = (b, c)$ and $t_{i+2} = (a, c)$. Without loss of generality the latter holds. Then, since $i, i+1 \in \mathrm{Des}(\phi(w))$,

$$D_iD_{i+1}D_i(w) = D_i(t_1, \ldots, t_{i-1}, (b, c), (a, c), t_i, t_{i+1}, \ldots, t_{n-1})$$

$$= (t_1, \ldots, t_{i-1}, (a, b), (b, c), t_i, t_{i+3}, \ldots, t_{n-1}).$$

On the other hand,

$$D_{i+1}D_iD_{i+1}(w) = D_{i+1}D_i(t_1, \ldots, t_{i-1}, t_i, (a, b), (b, c), t_{i+3}, \ldots, t_{n-1})$$

$$= (t_1, \ldots, t_{i-1}, (a, b), (b, c), t_i, t_{i+3}, \ldots, t_{n-1}).$$

Similarly, if t_{i+2} commutes with t_i and t_{i+1} then, since $i, i+1 \in \mathrm{Des}(\phi(w))$, letting $\mathrm{Supp}_{\{i,i+1\}} = \{a, b, c\}$ with $a < b < c$,

$$D_iD_{i+1}D_i(w) = D_{i+1}D_iD_{i+1}(w) = (t_1, \ldots, t_{i-1}, t_{i+2}, (a, b), (b, c), t_{i+3}, \ldots, t_{n-1}).$$

If t_{i+1} commutes with t_i and t_{i+2} then, since $i, i+1 \in \mathrm{Des}(\phi(w))$, letting $\mathrm{Supp}_{\{i,i+2\}} = \{a, b, c\}$ with $a < b < c$,

$$D_i D_{i+1} D_i(w) = D_{i+1} D_i D_{i+1}(w) = (t_1, \ldots, t_{i-1}, (a,b), t_{i+1}, (b,c), t_{i+3}, \ldots, t_{n-1}).$$

Case 3. $|\mathrm{Supp}_{\{i,i+1,i+2\}}(w)| = 4$.

Let $\mathrm{Supp}_{\{i,i+1,i+2\}}(w) = \{1, 2, 3, 4\}$ with $1 < 2 < 3 < 4$. By Proposition 3.5, there are exactly 16 possible options for (t_i, t_{i+1}, t_{i+2}), which are drawn in Figure 2.2. Looking at this figure, one can verify that the braid relation holds. □

5.2. **Maximal intervals in the weak order on F_n.**

Definition 5.5. For a permutation $\pi \in S_{n-1}$ let $s_{i_1} \cdots s_{i_k}$ be a reduced word for π in the alphabet of simple reflections. Define the operator $D_\pi : F_n \longrightarrow F_n$ by

$$D_\pi := D_{i_1} \cdots D_{i_k}.$$

Claim 5.6. The operator D_π is well-defined.

Proof. Since any two reduced words for the same permutation are connected by a sequence of braid relations, the result follows from Lemma 5.4. □

Lemma 5.7. *For every element $w \in F_n$, the lower interval $[e, w]$ is exactly $\{D_\pi(w) : \pi \in S_{n-1}\}$.*

Proof. By definition, $D_i(w) \le w$ for any $1 \le i \le n-2$ and therefore $D_\pi(w) \le w$ for any $\pi \in S_{n-1}$.

In the other direction, if $v \le w$ then v belongs to a geodesic from w to e in $G_T(n)$, part of which is a saturated chain $w = w_0 > w_1 > \ldots > w_k = v$ in $\mathrm{Weak}(F_n)$. Clearly each step is $w_j = D_{i_j}(w_{j-1})$ for some i_j $(1 \le j \le k)$. If the word $s_{i_k} \cdots s_{i_1}$ is not reduced then it can be brought, by relations as in Lemma 5.4, to a reduced word for some π, so that $v = D_\pi(w)$. □

Let $\pi_0 = [n-1, n-2, \ldots, 1]$ be the maximal element in the weak order on S_{n-1}.

Lemma 5.8. *For every maximal element w_0 in $\mathrm{Weak}(F_n)$ and every $\pi \in S_{n-1}$,*

$$\phi(D_\pi(w_0)) = \pi_0 \pi^{-1}.$$

Proof. By induction on $\mathrm{inv}(\phi(\pi))$. If $\pi = id$ then necessarily $\phi(D_\pi(w_0)) = \phi(w_0) = \pi_0$, the maximal element in $\mathrm{Weak}(S_{n-1})$, since otherwise there exists $1 \le j < n-1$ such that $\mathrm{inv}(\phi(w_0)s_j) > \mathrm{inv}(\phi(w_0))$. Then, by Lemma 3.10, $v = R_j(w_0)$ or $v = L_j(w_0)$ with $\mathrm{rank}(v) > \mathrm{rank}(w_0)$; thus, by Lemma 4.3, w_0 is not maximal.

For any $\pi \neq id$ there exist j such that $\pi = s_j\sigma$ with $\mathrm{inv}(\sigma) = \mathrm{inv}(\pi) - 1$. Then $\pi_0\pi^{-1} = \pi_0\sigma^{-1}s_j$ with $j \in \mathrm{Des}(\pi_0\sigma^{-1})$. By induction hypothesis $\phi(D_\sigma(w_0)) = \pi_0\sigma^{-1}$, thus $j \in \mathrm{Des}(\phi(D_\sigma(w_0)))$. Combining this with Observation 5.3 we obtain

$$\phi(D_\pi(w_0)) = \phi(D_j D_\sigma(w_0)) = \phi(D_\sigma(w_0))s_j = \pi_0\sigma^{-1}s_j = \pi_0\pi^{-1}.$$

$\square$

We conclude

Theorem 5.9. *Each maximal intervals in* Weak(F_n) *are isomorphic, as a poset, to the symmetric group* S_{n-1} *with the weak order.*

Proof. By Lemma 5.7 together with Lemma 5.8, ϕ, even when restricted to a maximal interval $[e, w_0]$, is onto S_{n-1}.

To prove that ϕ, when restricted to $[e, w_0]$, is one-to-one, it suffices to show that the cardinality of the interval $[e, w_0]$ is not bigger than the cardinality of the image S_{n-1}. But this follows from Lemma 5.4.

To prove that ϕ is a poset isomorphism, notice that since $\mathrm{inv}(\phi(w)s_j) < \mathrm{inv}(\phi(w))$ implies either $R_j(w) < w$ or $L_j(w) < w$,
w covers v in Weak(F_n)

$$\iff \quad v \in \{R_j(u), L_j(w) : \ 1 \le j < n-1\} \ \text{ and } \ \mathrm{inv}(\phi(w)) = \mathrm{inv}(\phi(v)) + 1,$$

$$\iff \quad \phi(v) \in \{\phi(w)s_j : \ 1 \le j < n-1\} \quad \text{and} \quad \mathrm{inv}(\phi(w)) = \mathrm{inv}(\phi(v)) + 1,$$

that is, $\phi(w)$ covers $\phi(v)$ in Weak(S_{n-1}).

$\square$

We can now summarize.

Theorem 5.10.

1. *The zero-Hecke algebra* $\mathcal{H}_{n-1}(0)$ *acts faithfully on* Weak(F_n).
2. *Lower intervals are invariant under this action.*

Proof. By Lemma 5.4, the down operators $\{D_i : \ 1 \le i < n-1\}$ respect the defining relations. Furthermore, by Lemma 5.7, lower intervals are invariant under these operators. By Theorem 5.9, the size of a maximal lower interval is equal to the dimension of the zero-Hecke algebra, thus the action is faithful.

$\square$

6. Number of maximal elements

In this section we prove

Theorem 6.1. *The number of maximal elements in* Weak(F_n) *is the Catalan number*

$$C_{n-1} = \frac{1}{n}\binom{2n-2}{n-1}.$$

For a q-analogue of this theorem see Corollary 8.5 below.

Recall the map $\phi : F_n \to S_{n-1}$ defined in Section 3. For $\pi \in S_{n-1}$, let

$$N(\pi) := \left|\{w \in F_n \mid \phi(w) = \pi^{-1}\}\right|.$$

For a subset $S = \{i_1, \ldots, i_k\}$ of $\{1, \ldots, n-1\}$, with $i_1 < \ldots < i_k$, let $\pi|_S$ be the subsequence $(\pi(i_1), \ldots, \pi(i_k))$, and let $p(\pi|_S)$ be the corresponding pattern (permutation) in the region $(1, \ldots, k)$, namely: the sequence obtained from $\pi|_S$ by writing 1 instead of the smallest letter, 2 instead of the second smallest letter, etc. For example, if $\pi = 3572146 \in S_7$ and $S = \{1, 2, 3, 7\}$ then $\pi|_S = 3576$ and $p(\pi|_S) = 1243$.

Also, for $1 \le i < j \le n$, let

$$A(i,j) := \{i, \ldots, j-1\} \setminus \{j-1\}$$

(of size $j - i - 1 \ge 0$) and

$$B(i,j) := \{j, \ldots, n, 1, \ldots, i-1\} \setminus \{n\}$$

(of size $n - j + i - 1 \ge 0$). Thus $A(i,j) \cup B(i,j) = \{1, \ldots, n\} \setminus \{j-1, n\}$, a disjoint union.

Lemma 6.2. *Let $\pi \in S_{n-1}$ and denote $j := 1 + \pi^{-1}(1)$. Then*

$$N(\pi) = \sum_{1 \le i < j} N(p(\pi|_{A(i,j)})) \cdot N(p(\pi|_{B(i,j)})).$$

Example 6.3. For $\pi = 25314 \in S_5$, $\pi^{-1} = 41352$ and $j - 1 = \pi^{-1}(1) = 4$. Thus, summing over $1 \le i \le 4$:

$$\begin{aligned} N(25314) &= N(p(253))N(p(4)) + N(p(53))N(p(24)) + N(p(3))N(p(254)) + \\ &\quad N(p())N(p(2534)) \\ &= N(132)N(1) + N(21)N(12) + N(1)N(132) + N()N(1423). \end{aligned}$$

Proof. Assume that $w \in F_n$ satisfies $\phi(w)(1) = j-1$. Then the first transposition in w has the form $t_1 = (i,j)$ for some $i < j$, and the corresponding partial product $\sigma_2 = t_2 \cdots t_{n-1} = (i, i+1, \ldots, j-1)(j, j+1, \ldots, n, 1, \ldots, i-1)$. This follows from the fact that for any $1 \le k \le n-1$, the transition from any σ_{k+1} to σ_k amounts to “merging” two of the cycles of σ_{k+1} into one cycle, in which the original cycles form complementary cyclic intervals. It follows that $(i, i+1, \ldots, j-1)$ is the product of those t_k with $k > 1$ which permute two elements of the set $\{i, i+1, \ldots, j-1\}$, and thus have $\phi(w)(k)$ in this set (but of course $\phi(w)(k) \ne \phi(w)(1) = j-1$); and similarly $(j, j+1, \ldots, n, 1, \ldots, i-1)$ is the product of those t_k with $k > 1$ which permute two elements of the set $\{j, j+1, \ldots, n, 1, \ldots, i-1\}$, and thus have $\phi(w)(k)$ in this set (but of course $\phi(w)(k) \ne n$). w is a shuffle of these two sequences of transpositions, with $t_1 = (i,j)$ added as a prefix. The ϕ values at these two sequences compose the ranges $A(i,j)$ and $B(i,j)$, respectively. When they are mapped to the ranges

$\{1, \ldots, j-i-1\}$ and $\{1, \ldots, n-j+i-1\}$ as above, letting $\pi := \phi(w)^{-1}$ we get the corresponding permutations $p(\pi|_{A(i,j)})$ and $p(\pi|_{B(i,j)})$.

□

Proof of Theorem 6.1. Let π be the longest element in S_{n-1}, i.e. $\pi(i) = n - i$ $(1 \le i \le n-1)$. Of course, $\pi^{-1} = \pi$. Denote the corresponding $N(\pi)$ by N_n. If $w \in F_n$ has $\phi(w) = \pi$ then $j = n$ and therefore, for each $1 \le i \le n-1$, $A(i,j) = \{i, \ldots, n-2\}$ and $B(i,j) = \{1, \ldots, i-1\}$. Thus $\pi|_{A(i,j)}$ and $\pi|_{B(i,j)}$ are decreasing sequences, and the corresponding permutations $p(\pi|_{A(i,j)})$ and $p(\pi|_{B(i,j)})$ are the longest elements in S_{n-i-1} and S_{i-1}, respectively. It follows, by Lemma 6.2, that

$$N_n = \sum_{i=1}^{n-1} N_{n-i} N_i \qquad (\forall n \ge 2)$$

which is the recursion formula of the Catalan numbers C_{n-1}. Together with the initial value $N_1 = 1 = C_0$ this implies that

$$N_n = C_{n-1} = \frac{1}{n}\binom{2n-2}{n-1},$$

as claimed

□

The following lemma will be used later.

Lemma 6.4. *Each maximal element in* $\mathrm{Weak}(F_n)$ *contains the factor* $(1, n)$.

Proof. By induction on n. Obviously, the lemma holds for $n = 2$. Assume that it holds for every $m < n$. By Observation 3.3, any word $v = (t_1, \ldots, t_{n-1}) \in F_n$ with $\phi(v) = [n, n-1, \ldots, 1]$ must end with $t_{n-1} = (1, j)$ for some $1 < j \le n$. By same arguments as in the proof of Theorem 6.1, $(t_1, \ldots, t_{n-2})$ is a shuffle of a reduced word of an element $u \in F_{j-1}$ whose image is $[j, j-1, \ldots, 2]$ and an element $w \in F_{n-j+1}$ whose image is $[n, n-1, \ldots, j+1, 1]$. By induction hypothesis, w contains the factor $(1, n)$.

□

7. Right and left inversions

Recall that the weak order on S_n is characterized by the inversion set, namely: $\pi \le \sigma$ in the (right) weak order on S_n if and only if $\mathrm{Inv}(\pi) \subseteq \mathrm{Inv}(\sigma)$, where $\mathrm{Inv}(\pi) := \{(i,j) : \ i < j \text{ and } \pi^{-1}(i) > \pi^{-1}(j)\}$ (see, e.g., [4, Cor. 1.5.2, Prop. 3.1.3]).

The goal of this section is to provide a similar characterization for $\mathrm{Weak}(F_n)$.

Definition 7.1. The set of *inversions* of $w = (t_1, \ldots, t_{n-1}) \in F_n$ is $\mathrm{Inv}(w) := \mathrm{Inv}(\phi(w))$. A pair $(i,j) \in \mathrm{Inv}(\phi(w))$ is a *right inversion* if there exist $a < b < c$ such that $t_{\phi(w)^{-1}(j)} = (a,c)$ and $t_{\phi(w)^{-1}(i)} = (a,b)$, or there exist $a < b < c < d$ such that $t_{\phi(w)^{-1}(j)} = (a,d)$ and $t_{\phi(w)^{-1}(i)} = (b,c)$;

a *left inversion* if there exist $a < b < c$ such that $t_{\phi(w)^{-1}(j)} = (b,c)$ and $t_{\phi(w)^{-1}(i)} = (a,c)$, or there exist $a < b < c < d$ such that $t_{\phi(w)^{-1}(j)} = (b,c)$ and $t_{\phi(w)^{-1}(i)} = (a,d)$;

a *neutral inversion* if there exist $a < b < c < d$ such that $t_{\phi(w)^{-1}(j)} = (c,d)$ and $t_{\phi(w)^{-1}(i)} = (a,b)$.

Remark 7.2. If $t_{\phi(w)^{-1}(i)} = (a,b)$, $a < b$, write I_i for the interval $[a,b) \subseteq [1,n]$. Then $(i,j) \in \mathrm{Inv}(\phi(w))$ is

a right inversion if and only if $I_i \subset I_j$;
a left inversion if and only if $I_j \subset I_i$;
a neutral inversion if and only if $I_i \cap I_j = \emptyset$.

Denote the set of right, left and neutral inversions of $w \in F_n$ by $\mathrm{Inv}_R(w)$, $\mathrm{Inv}_L(w)$ and $\mathrm{Inv}_N(w)$, respectively. Denote also $\mathrm{Inv}_{LN}(w) := \mathrm{Inv}_L(w) \cup \mathrm{Inv}_N(w)$ and $\mathrm{Inv}_{RN}(w) := \mathrm{Inv}_R(w) \cup \mathrm{Inv}_N(w)$.

By Lemma 3.9, for every $w \in F_n$,

$$\mathrm{Inv}(w) = \mathrm{Inv}_R(w) \cup \mathrm{Inv}_L(w) \cup \mathrm{Inv}_N(w), \tag{10}$$

a disjoint union.

Example 7.3. For $w = ((3,4),(1,5),(5,6),(1,4),(1,2)) \in F_6$, $\phi(w) = [3,4,5,2,1]$. Then $\mathrm{Inv}(w) = \{(1,2),(1,3),(1,4),(1,5),(2,3),(2,4),(2,5)\}$ is a disjoint union of $\mathrm{Inv}_R(w) = \{(1,2),(1,4),(2,5)\}$, $\mathrm{Inv}_N(w) = \{(1,3),(1,5),(2,5)\}$ and $\mathrm{Inv}_L(w) = \{(2,3)\}$.

Define the involution $\iota : F_n \longrightarrow F_n$ by $[\iota(t_1,\ldots,t_{n-1}) := (t_{n-1}^{w_0},\ldots,t_1^{w_0}),]$ where $g^h := h^{-1}gh$ and $w_0 := [n-1,\ldots,1]$ is the unique maximal element in $\mathrm{Weak}(S_{n-1})$.

Observation 7.4.

1. For every $w \in F_n$, $\phi(\iota(w)) = \phi(w)^{w_0}$, $\mathrm{Inv}_N(\iota(w)) = \mathrm{Inv}_N(w)^{w_0}$, $\mathrm{Inv}_R(\iota(w)) = \mathrm{Inv}_L(w)^{w_0}$ and $\mathrm{Inv}_L(\iota(w)) = \mathrm{Inv}_R(w)^{w_0}$.
2. ι is a rank preserving automorphism of $\mathrm{Weak}(F_n)$.

Proposition 7.5.
Each of the pairs $(\mathrm{Inv}_L(u), \mathrm{Inv}_{RN}(u))$ *and* $(\mathrm{Inv}_R(u), \mathrm{Inv}_{LN}(u))$ *determines the word* u. *Namely, for any pair of words* $u, v \in F_n$

$$\mathrm{Inv}_L(u) = \mathrm{Inv}_L(v) \quad \text{and} \quad \mathrm{Inv}_{RN}(u) = \mathrm{Inv}_{RN}(v) \quad \Longrightarrow \qquad u = v, \tag{11}$$

and

$$\text{(12)} \qquad \mathrm{Inv}_R(u) = \mathrm{Inv}_R(v) \ \text{ and } \ \mathrm{Inv}_{LN}(u) = \mathrm{Inv}_{LN}(v) \quad \Longrightarrow \quad u = v.$$

Proof. To prove (11), it suffices to show that $\mathrm{Inv}_L(u)$ and $\mathrm{Inv}(u) = \mathrm{Inv}_{RN}(u) \cup \mathrm{Inv}_L(u)$ determine u. Let $u = (t_1, \ldots, t_{n-1})$. First, we will show that $\mathrm{Inv}(u)$ and $\mathrm{Inv}_L(u)$ determine t_{n-1}, the last transposition in u. Assume that $t_{n-1} = (i,j)$ with $i < j$. Clearly, $\mathrm{Inv}(u) = \mathrm{Inv}(\phi(u))$ determines $\phi(u) \in S_{n-1}$. By definition of ϕ, $i = \phi(u)(n-1)$. To recover j, we will show that there are exactly $j-i-1$ left inversions involving $i = \phi(u)(n-1)$; namely, the cardinality of the set $\{(i,k) : 1 \le k \le n-1\} \cap \mathrm{Inv}_L(u)$ is exactly $j-i-1$. Indeed, by Remark 7.2, left inversions of $\phi(u)$ involving i correspond bijectively to edges of $G(u)$ with both endpoints in the interval $I_i = [i,j]$, except the edhe (i,j) itself. By Proposition 3.5(i), there are exactly $j-i-1$ such edges.

Now, for every $1 \le k \le n-2$, we will recover t_k in a similar fashion. Let $t_k = (i,j)$. Then $i = \sigma_{k+1}(\phi(u)(k))$. Also, $t_1 \cdots t_k = c\sigma_{k+1}^{-1}$ is a product of $n-k$ disjoint cycles. Let $\tilde{c}$ be the cycle containing i. Then, by arguments as above

$$\#\{(i,r) : \ r \in \tilde{c}\} \cap \sigma_{k+1}(\mathrm{Inv}_L(u)) = j - i - 1.$$

To prove (12), notice that by Observation 7.4.1, $\mathrm{Inv}(u)$ and $\mathrm{Inv}_L(u)$ determine u if and only if $\mathrm{Inv}(u)$ and $\mathrm{Inv}_R(u)$ determine u.

□

Furthermore, the following strengthening of Proposition 7.5 holds.

Proposition 7.6. *For $u, v \in F_n$,*

$$u \le v \text{ in } \mathrm{Weak}(F_n) \iff \mathrm{Inv}_R(u) \subseteq \mathrm{Inv}_R(v) \text{ and } \mathrm{Inv}_{LN}(u) \subseteq \mathrm{Inv}_{LN}(v).$$

Proof. First, we prove that the inclusions $\mathrm{Inv}_R(u) \subseteq \mathrm{Inv}_R(v)$ and $\mathrm{Inv}_{LN}(u) \subseteq \mathrm{Inv}_{LN}(v)$ imply $u \le v$. This will be proved by induction on the difference $\mathrm{rank}(v) - \mathrm{rank}(u)$.

If $\mathrm{rank}(v) = \mathrm{rank}(u)$ then, by Lemma 4.3, $\mathrm{inv}(\phi(u)) = \mathrm{inv}(\phi(v))$. Hence $|\mathrm{Inv}(u)| = |\mathrm{Inv}(v)|$. Furthermore by (10),

$$|\mathrm{Inv}_R(u)| + |\mathrm{Inv}_{LN}(u)| = |\mathrm{Inv}(u)| = |\mathrm{Inv}(v)| = |\mathrm{Inv}_R(v)| + |\mathrm{Inv}_{LN}(v)|.$$

Combining this together with the assumed inclusions implies that $\mathrm{Inv}_{LN}(u) = \mathrm{Inv}_{LN}(v)$, $\mathrm{Inv}_R(u) = \mathrm{Inv}_R(v)$. Hence, by Proposition 7.5, the equality $u = v$ holds.

If $\mathrm{rank}(u) < \mathrm{rank}(v)$ then, by Lemma 4.3, $\mathrm{inv}(\phi(u)) < \mathrm{inv}(\phi(v))$. By the assumed inclusions together with (10), $\mathrm{Inv}(u) \subseteq \mathrm{Inv}(v)$. Hence $\phi(u) < \phi(v)$ in $\mathrm{Weak}(S_{n-1})$, thus there exists $1 \le i < n-1$, such that $\phi(u) \le \phi(v)s_i < \phi(v)$. By the proof of Theorem 5.9, either $L_i v < v$ and $R_i v \le v$, or $L_i v = v$ and $R_i v < v$. Assume that $L_i v = v$ and $R_i v < v$. Then, since $\phi(u) \le \phi(v)s_i < \phi(v)$, $\mathrm{Inv}(v) \setminus \mathrm{Inv}(R_i v) = \{s_i^{\phi(v)}\}$, $s_i^{\phi(v)} \in \mathrm{Inv}_R(v)$ and $s_i^{\phi(v)} \notin \mathrm{Inv}(u)$. This together with the assumed inclusions implies that $\mathrm{Inv}_R(u) \subseteq \mathrm{Inv}_R(R_i v)$ and

$\mathrm{Inv}_{LN}(u) \subseteq \mathrm{Inv}_{LN}(R_i v)$. By the induction hypothesis $u \leq R_i v$ in $\mathrm{Weak}(F_n)$, thus by transitivity $u \leq v$. The analysis of the case $L_i v < v$ and $R_i v = v$ is similar.

If $R_v < v$ and $L_i v < v$ then , by Lemma 4.5, $R_i v = L_i v$. Then the i-th and the $i+1$-st transpositions in v, t_i and t_{i+1}, commute. Also $\phi(v)(i) > \phi(v)(i+1)$. There are three options: $t_i = (c,d)$ and $t_{i+1} = (a,b)$, or $t_i = (a,d)$ and $t_{i+1} = (b,c)$, or $t_i = (b,c)$ and $t_{i+1} = (a,d)$, for some $1 \leq a < b < c < d \leq n$. In first case, $\mathrm{Inv}_R(v) = \mathrm{Inv}_R(R_i v)$, $\mathrm{Inv}_L(v) = \mathrm{Inv}_L(R_i v)$ and $\mathrm{Inv}_N(v) = \mathrm{Inv}_N(R_i v) \cup \{s_i^{\phi(u)}\}$. In the second case, $\mathrm{Inv}_R(v) = \mathrm{Inv}_R(R_i v) \cup \{s_i^{\phi(u)}\}$, $\mathrm{Inv}_L(v) = \mathrm{Inv}_L(R_i v)$ and $\mathrm{Inv}_N(v) = \mathrm{Inv}_N(R_i v)$. In last case, $\mathrm{Inv}_L(v) = \mathrm{Inv}_L(R_i v) \cup \{s_i^{\phi(u)}\}$, $\mathrm{Inv}_R(v) = \mathrm{Inv}_R(R_i v)$ and $\mathrm{Inv}_N(v) = \mathrm{Inv}_N(R_i v)$. In all three cases, since $s_i^{\phi(v)} \notin \mathrm{Inv}(u)$, $\mathrm{Inv}_R(u) \subseteq \mathrm{Inv}_R(R_i v)$ and $\mathrm{Inv}_{LN}(u) \subseteq \mathrm{Inv}_{LN}(R_i v)$. Thus, by the induction hypothesis $u \leq R_i v$, and by transitivity $u \leq v$.

In the opposite direction, to prove that $u \leq v$ implies $\mathrm{Inv}_R(u) \subseteq \mathrm{Inv}_R(v)$ and $\mathrm{Inv}_{LN}(u) \subseteq \mathrm{Inv}_{LN}(v)$, it should be noticed that, by transitivity, it suffices to verify that these inclusions hold on covering relations. This may be done by a case by case elementary analysis, noting that if v covers u in $\mathrm{Weak}(F_n)$ then there exist $1 \leq i < n-1$ such that $\mathrm{Inv}(v) = \mathrm{Inv}(u) \cup \{s_i^{\phi(u)}\}$, and either $v = R_i u \neq L_i u$ or $v = L_i u \neq R_i u$ or $v = R_i u = L_i u$. In first case, $\mathrm{Inv}_R(u) = \mathrm{Inv}_R(v) \cup \{s_i^{\phi(u)}\}$, $\mathrm{Inv}_L(u) = \mathrm{Inv}_L(v)$ and $\mathrm{Inv}_N(v) = \mathrm{Inv}_N(u)$. In second case, $\mathrm{Inv}_R(u) = \mathrm{Inv}_R(v)$, $\mathrm{Inv}_L(u) = \mathrm{Inv}_L(v) \cup \{s_i^{\phi(u)}\}$ and $\mathrm{Inv}_N(v) = \mathrm{Inv}_N(u)$. Finally, if $v = R_i u = L_i u$ then letting t_i and t_{i+1} be the i-th and $i+1$-st transpositions in v either $t_i = (c,d)$ and $t_{i+1} = (a,b)$; or $t_i = (a,d)$ and $t_{i+1} = (b,c)$; or $t_i = (b,c)$ and $t_{i+1} = (a,d)$, for some $1 \leq a < b < c < d \leq n$. Then either $\mathrm{Inv}_R(v) = \mathrm{Inv}_R(R_i v)$, $\mathrm{Inv}_L(v) = \mathrm{Inv}_L(R_i v)$ and $\mathrm{Inv}_N(v) = \mathrm{Inv}_N(R_i v) \cup \{s_i^{\phi(u)}\}$; or: $\mathrm{Inv}_R(v) = \mathrm{Inv}_R(R_i v) \cup \{s_i^{\phi(u)}\}$, $\mathrm{Inv}_L(v) = \mathrm{Inv}_L(R_i v)$ and $\mathrm{Inv}_N(v) = \mathrm{Inv}_N(R_i v)$; or: $\mathrm{Inv}_L(v) = \mathrm{Inv}_L(R_i v) \cup \{s_i^{\phi(u)}\}$, $\mathrm{Inv}_R(v) = \mathrm{Inv}_R(R_i v)$ and $\mathrm{Inv}_N(v) = \mathrm{Inv}_N(R_i v)$; respectively.

□

Corollary 7.7. *For any pair* $u, v \in F_n$,

$$\begin{aligned} &\mathrm{Inv}_R(u) \subseteq \mathrm{Inv}_R(v) \quad \textit{and} \quad \mathrm{Inv}_{LN}(u) \subseteq \mathrm{Inv}_{LN}(v) \\ \iff &\mathrm{Inv}_L(u) \subseteq \mathrm{Inv}_L(v) \quad \textit{and} \quad \mathrm{Inv}_{RN}(u) \subseteq \mathrm{Inv}_{RN}(v). \end{aligned}$$

Proof. By Observation 7.4, one can replace the roles of right inversions and left inversions in Proposition 7.6. Thus, for any pair $u, v \in F_n$, $u \leq v$ in $\mathrm{Weak}(F_n)$ if and only if the following inclusions hold: $\mathrm{Inv}_L(u) \subseteq \mathrm{Inv}_L(v)$ and $\mathrm{Inv}_{RN}(u) \subseteq \mathrm{Inv}_{RN}(v)$. Combining this with Proposition 7.6 completes the proof.

□

8. Right inversions and q-Catalan numbers

A q-analogue of Theorem 6.1 is given in this section.

8.1. **Enumeration by right and left inversions.** Carlitz and Riordan [6] defined a q-Catalan number $C_n(q)$ using the recursion

$$C_{n+1}(q) := \sum_{k=0}^{n} q^{(k+1)(n-k)} C_k(q) C_{n-k}(q) \qquad (n \ge 0)$$

with $C_0(q) := 1$. For combinatorial interpretations of this number and further information see e.g. [8, 5, 15, 16]; in particular, $C_n(q)$ counts Dyck paths of length n with respect to the area above the path; see, e.g., [8].

Definition 8.1. Define (q,t)-*Catalan numbers* by

$$\tilde{C}_{n+1}(q,t) := \sum_{k=0}^{n} q^k t^{n-k} \tilde{C}_k(q,t) \tilde{C}_{n-k}(q,t) \qquad (n \ge 0)$$

with

$$\tilde{C}_0(q,t) = 1.$$

Observation 8.2. For every $n \ge 0$,

$$\tilde{C}_n(q,t) = \tilde{C}_n(t,q) \tag{13}$$

and

$$C_n(q) = q^{\binom{n}{2}} \tilde{C}_n(q^{-1},1). \tag{14}$$

Definition 8.3. For $w \in F_n$, denote the cardinality of $\mathrm{Inv}_R(w)$ by $\mathrm{inv}_R(w)$ and the cardinality of $\mathrm{Inv}_L(w)$ by $\mathrm{inv}_L(w)$.

Proposition 8.4. *For every positive integer n,*

$$\sum_{w \in F_{n+1},\ \mathrm{rank}(w) = \binom{n}{2}} q^{\mathrm{inv}_R(w)} t^{\mathrm{inv}_L(w)} = \tilde{C}_n(q,t). \tag{15}$$

Proof. By induction on n. For $n = 1$, equation (15) obviously holds.

For the induction step recall that, by Lemma 6.4, any maximal element in w in $\mathrm{Weak}(F_{n+1})$ contains the factor $(1, n+1)$. Denote by $F_{n+1,k}$ the set of maximal elements in $\mathrm{Weak}(F_{n+1})$ whose $(k+1)$-st factor is $(1, n+1)$. By the proof of Theorem 6.1, this set is characterized as follows: $w = (t_1, \ldots, t_n) \in F_{n+1,k}$ if and only if $t_1 t_2 \cdots t_k$ is a reduced word for the cycle $(n-k, \ldots, n)$, where $\phi(t_1 t_2 \cdots t_k) = [n-1, \ldots, n-k]$, and $t_{k+2} t_{k+3} \cdots t_n$ is a reduced word for the cycle $(1, 2, \ldots, n-k)$, where $\phi(t_{k+2} t_{k+3} \cdots t_n) = [n-k-1, n-k-2, \ldots, 1]$.

For every $w = t_1 \cdots t_n \in F_{n+1,k}$ the following holds: for every $1 \le i \le k$ there exist $n-k \le c < d \le n$ such that $t_i = (c,d)$; for every $k+1 < j \le n$ there exist $1 \le a < b < n-k$ such that $t_j = (a,b)$. Thus, by Definition 7.1, all pairs (t_i, t_j), with $i \le k$ and $j > k+1$, are neutral inversions; all pairs (t_i, t_{k+1}), $1 \le i \le k$, are left inversions and all pairs (t_{k+1}, t_j), $k+1 < j \le n$, are right inversions.

This implies that for every $0 \le k \le n$

$$\sum_{\{w\in F_{n+1,k}:\ \mathrm{rank}(w)=\binom{n}{2}\}} q^{\mathrm{inv}_R(w)} t^{\mathrm{inv}_L(w)}$$

$$= q^{n-k-1}t^k \cdot \sum_{\{w\in F_{k+1}:\ \mathrm{rank}(w)=\binom{k}{2}\}} q^{\mathrm{inv}_R(w)} t^{\mathrm{inv}_L(w)} \cdot \sum_{\{w\in F_{n-k+1}:\ \mathrm{rank}(w)=\binom{n-k}{2}\}} q^{\mathrm{inv}_R(w)} t^{\mathrm{inv}_L(w)}$$

$$= q^{n-k-1}t^k \cdot \tilde{C}_k(q,t) \cdot \tilde{C}_{n-k}(q,t).$$

The last equality follows from the induction hypothesis. Thus

$$\sum_{\{w\in F_{n+1}:\ \mathrm{rank}(w)=\binom{n}{2}\}} q^{\mathrm{inv}_R(w)} t^{\mathrm{inv}_L(w)} = \sum_{k=0}^{n} \sum_{\{w\in F_{n+1,k}:\ \mathrm{rank}(w)=\binom{n}{2}\}} q^{\mathrm{inv}_R(w)} t^{\mathrm{inv}_L(w)}$$

$$= \sum_{k=0}^{n} q^{n-k}t^k \cdot \tilde{C}_k(q,t) \cdot \tilde{C}_{n-k}(q,t) = \tilde{C}_{n+1}(q,t).$$

The last equality follows from the defining recursion of $\tilde{C}_n(q,t)$ together with (13).

□

Corollary 8.5. *For every positive integer n,*

$$\sum_{w\in F_{n+1},\ \mathrm{rank}(w)=\binom{n}{2}} q^{\mathrm{inv}_R(w)} = q^{\binom{n}{2}} C_n(q^{-1})$$

or equivalently

$$\sum_{w\in F_{n+1},\ \mathrm{rank}(w)=\binom{n}{2}} q^{\mathrm{inv}_{LN}(w)} = C_n(q).$$

Proof. Proposition 8.4 together with (14). □

8.2. **Another combinatorial interpretation.**

Consider the set $D(n)$ of Dyck paths of length $2n$. Each $p \in D(n)$ is a sequence $(\epsilon_1, \ldots, \epsilon_{2n})$, with

$$\epsilon_i \in \{1, -1\} \qquad (\forall i),$$

such that

$$\epsilon_1 + \ldots + \epsilon_{2n} = 0$$

and

$$\epsilon_1 + \ldots + \epsilon_k \ge 0 \qquad (\forall k).$$

Geometrically, p corresponds to a lattice path from $(0,0)$ to $(2n,0)$, with admissible steps $(1,1)$ and $(1,-1)$, which never goes below the horizontal line (*baseline*, in the sequel) connecting its endpoints.

Definition 8.6. With each $p \in D(n)$ associate the following two statistics:

$$\begin{aligned}
\mathrm{area}(p) &:= \binom{n}{2} - \#\{(i,j) \,|\, 1 \le i < j \le 2n,\, -\epsilon_i = \epsilon_j = 1\} \\
&= -\binom{n+1}{2} + \#\{(i,j) \,|\, 1 \le i < j \le 2n,\, \epsilon_i = -\epsilon_j = 1\}, \\
\mathrm{bmaj}(p) &:= \sum_{i \,:\, -\epsilon_i = \epsilon_{i+1} = 1} \max\{k \,|\, (\forall 1 \le t \le 2k)\, \epsilon_{i+1} + \ldots + \epsilon_{i+t} \ge 0\}.
\end{aligned}$$

Geometrically, $\mathrm{area}(p)$ is the area between the path p and the "horizontal" path $p_0 = (1, -1, 1, -1, ...)$, where the square with vertices $(0,0)$, $(1,1)$, $(2,0)$ and $(1,-1)$ is considered to have area 1; while $\mathrm{bmaj}(p)$ is the sum, over all internal local minima of the path p, of half the length of the maximal horizontal segment starting at the minimum point and stretching to the right without crossing (but possibly touching) p. Note that $\mathrm{area}(p)$ is (a linear function of) the inversion number of the sequence p.

Proposition 8.7. *For every positive integer n*

$$\sum_{p \in D(n)} q^{\mathrm{area}(p)} t^{\mathrm{bmaj}(p)} = \tilde{C}_n(q,t).$$

Note that for $t = 1$ this is a classical result [8].

Proof. It suffices to show that the LHS, to be denoted $C'_n(q,t)$, satisfies the recursion in Definition 8.1; it clearly satisfies $C'_0(q,t) = 1$.

let $p \in D(n+1)$, and define $k := \min\{t \ge 0 \,|\, \epsilon_1 + \ldots + \epsilon_{2t+2} = 0\}$, so that $2k+2$ is the first time that p returns to the baseline. Clearly $0 \le k \le n$. We can write p as the concatenation of sequences

$$p = (1) * p' * (-1) * p'',$$

where $p' \in D(k)$ and $p'' \in D(n-k)$. Now clearly

$$\mathrm{area}(p) = k + \mathrm{area}(p') + \mathrm{area}(p'')$$

and

$$\mathrm{bmaj}(p) = (n-k) + \mathrm{bmaj}(p') + \mathrm{bmaj}(p''),$$

since the path p has an internal local minimum at $(2k+2, 0)$, in addition to the internal local minima of p' and p''. Thus

$$C'_{n+1}(q,t) = \sum_{k=0}^{n} q^k t^{n-k} C'_k(q,t) C'_{n-k}(q,t),$$

and this completes the proof. □

Corollary 8.8. area *and* bmaj *are equidistributed:*

$$\sum_{p \in D(n)} q^{\mathrm{area}(p)} = \sum_{p \in D(n)} q^{\mathrm{bmaj}(p)} = q^{\binom{n}{2}} C_n(q^{-1})$$

Proof. Proposition 8.7 together with both parts of Observation 8.2. ☐

Corollary 8.9.

$$\sum_{\{w\in F_{n+1}:\ \mathrm{rank}(w)=\binom{n}{2}\}} q^{\mathrm{inv}_R(w)}t^{\mathrm{inv}_L(w)} = \sum_{p\in D(n)} q^{\mathrm{area}(p)}t^{\mathrm{bmaj}(p)}.$$

Proof. Proposition 8.4 together with Proposition 8.7. ☐

9. Radius and diameter

Definition 9.1. Let $G=(V,E)$ be a finite connected undirected graph, and let $d(\cdot,\cdot)$ be the corresponding metric on its vertex set V. The *radius* of G is

$$\mathrm{rad}(G) := \min_{v\in V}\max_{w\in V} d(v,w)$$

and its *diameter* is

$$\mathrm{diam}(G) := \max_{v,w\in V} d(v,w).$$

Recall the Hurwitz graph $G_T(n)$ (Definition 2.2), and let $\mathrm{Cayley}(S_{n-1})$ be the Cayley graph of the group S_{n-1} with respect to its standard Coxeter generators.

Lemma 9.2. *The map* $\phi : G_T(n) \to \mathrm{Cayley}(S_{n-1})$ *is a contraction:*

$$d(\phi(v),\phi(w)) \le d(v,w) \qquad (\forall v,w\in F_n).$$

Proof. It suffices to show that if $d(v,w)=1$ then $d(\phi(v),\phi(w))\le 1$. Let $v,w\in F_n$ with $d(v,w)=1$ in $G_T(n)$. Thus either $v=R_j(w)$ or $v=L_j(w)$ for some j. If the edge connecting v and w in $G_T(n)$ is deleted during the passage to $\mathrm{Weak}(F_n)$ (see Corollary 4.4) then, by Lemma 3.10, $\phi(v)=\phi(w)$. Otherwise clearly $d(\phi(v),\phi(w))=1$. ☐

Theorem 9.3. *The radius of the Hurwitz graph*

$$\mathrm{rad}(G_T(n)) = \binom{n-1}{2}.$$

Proof. Since only edges connecting vertices of the same rank are deleted when passing from $G_T(n)$ to the Hasse diagram of $\mathrm{Weak}(F_n)$, it follows that the distance $d(e,w)$ from the unique minimal element e of $\mathrm{Weak}(F_n)$ to an arbitrary $w\in F_n$ is the same, whether measured in $G_T(n)$ or in the Hasse diagram. Thus, by Theorem 5.9 and a well-known property of S_{n-1},

$$\max_{w\in F_n} d(e,w) = \max_{\pi\in S_{n-1}} d(id,\pi) = \binom{n-1}{2}.$$

It follows that

$$\mathrm{rad}(G_T(n)) \le \binom{n-1}{2}.$$

Consider now an arbitrary $v \in F_n$. By the proof of Theorem 5.9, $\phi : F_n \to S_{n-1}$ is surjective. Thus there exists $w \in F_n$ with

$$\phi(w) = \phi(v) \cdot \pi_0,$$

where $\pi_0 = [n-1, \dots, 1] \in S_{n-1}$. For this w it follows from Lemma 9.2 that

$$d(v, w) \geq d(\phi(v), \phi(w)) = \binom{n-1}{2},$$

and therefore

$$\operatorname{rad}(G_T(n)) \geq \binom{n-1}{2}.$$

□

Clearly, for any graph G,

$$\operatorname{rad}(G) \leq \operatorname{diam}(G) \leq 2 \operatorname{rad}(G).$$

Corollary 9.4. *The diameter of $G_T(n)$ satisfies*

$$\frac{1}{2}(n-1)(n-2) \leq \operatorname{diam}(G_T(n)) \leq (n-1)(n-2).$$

The upper bound can be improved.

Proposition 9.5. *The diameter of $G_T(n)$ satisfies*

$$\frac{1}{2}(n-1)(n-2) \leq \operatorname{diam}(G_T(n)) \leq \frac{3}{4}(n-1)^2.$$

Proof. We will get from two given words in F_n same word by a careful use of the bubble sorting algorithm. While in classical bubble sorting one word is left unchanged, here the situation is a little different. The procedure in a general step is the following:

At the a general step, we are given w and v, which are two T-reduced words of a permutation $\pi \in S_n$. We equate first or last factor as follows: By Proposition 3.5(i), there is a letter $i \in \{1, \dots, n\}$, which appears only once in a word w. Without loss of generality, i appears in one of the first $\lceil \frac{\ell_T(\pi)}{2} \rceil$ factors of w, say in the j-th factor. We can push the appearance of i to the left using $j-1$ slides relations, keeping the number of the appearances of i to be one.

Now, we push the rightmost appearance of i in v to the left; if i does not appear in the left factor involving in the slide we push the appearance of i to the left; each time that the slide involves another appearance of i, we decrease the number of appearances of i by one; after at most $\ell(\pi) - 1$ slides we get a word with one appearance of i at the first factor. Thus, in both resulting words the letter i appears in the first factor only.

Now, observe that if the only appearance of a letter $i \in \{1, \dots, n\}$ in a T-reduced word of $\pi \in S_n$ is in the first factor then this factor is $(i, \pi(i))$ (similarly, if i appears in the last factor only then this factor is $(i, \pi^{-1}(i))$). This implies that the first factor in both resulting words is $(i, \pi(i))$. Omit first factor and operate same procedure of equating first or last letter in the resulting words, and so on.

Notice that at each step we reduce the T-length of the remaining word by one. Hence, altogether we used at most

$$\sum_{\ell=n-1}^{2} (\ell - 1 + \lfloor \ell/2 \rfloor) \leq \frac{3}{4}(n-1)^2.$$

slides to get from any two words in F_n identical words, completing the proof.

□

Conjecture 9.6. *The diameter of* $G_T(n)$ *is* $\binom{n-1}{2} + O(n)$.

Computer experiments for $n < 8$ seem to suggest that the diameter is actually $\left\lfloor \frac{(n-1)^2}{2} \right\rfloor - 1$.

Acknowledgements. Thanks to C. Athanasiadis, O. Bernardi, C. Keller, A. Kalka, A. Postnikov and V. Reiner for useful discussions, comments and references.

References

[1] D. Armstrong, *Generalized noncrossing partitions and combinatorics of Coxeter groups*, Mem. Amer. Math. Soc. **202** (2009), no. 949.

[2] M. Autord and P. Dehornoy, *On the distance between the expressions of a permutation*, `arXiv:0902.3074`.

[3] D. Bessis, *The dual braid monoid*, Ann. Sci. École Norm. Sup. (4) **36** (2003), 647–683.

[4] A. Björner and F. Brenti, Combinatorics of Coxeter groups, Graduate Texts in Mathematics 231, Springer, New York, 2005.

[5] L. M. Butler and W. P. Flanigan, *A note on log-convexity of q-Catalan numbers*, Ann. Comb. **11** (2007), 369–373.

[6] L. Carlitz and J. Riordan, *Two element lattice permutations and their q-generalization*, Duke J. Math. **31** (1964) 371–388.

[7] J. Dénes, *The representation of a permutation as the product of a minimal number of transpositions and its connection with the theory of graphs*, Publ. Math. Inst. Hungar. Acad. Sci. **4** (1959), 63–71.

[8] J. Fürlinger and J. Hofbauer, *q-Catalan numbers*, J. Combin. Theory Ser. A **40** (1985), 248–264.

[9] D. A. Gewurz and F. Merola, *Some factorisations counted by Catalan numbers*, European J. Combin. **27** (2006), 990–994.

[10] I. Goulden and A. Yong, *Tree-like properties of cycle factorizations*, J. Combin. Theory Ser. A **98** (2002), 106–117.

[11] C. Hernando, F. Hurtado, A. Márquez, M. Mora and M. Noy, *Geometric tree graphs of points in convex position*, Discrete Appl. Math. **93** (1999), 51–66.

[12] A. Hurwitz, *Ueber Riemann'sche Flächen mit gegebenen Verzweigungspunkten*, Math. Ann. **39** (1891), 1–61.

[13] G. Kreweras, *Sur les partitions non croisées d'un cycle*, Discrete Math. **1** (1972), 333–350.

[14] V. Reiner and Y. Roichman, *Diameter of reduced words*, Trans. Amer. Math. Soc., to appear; `arXiv:0906.4768`.

[15] B. Sagan, and C. Savage, *Mahonian pairs*, `arXiv:1101.4332`.

[16] N. J. A. Sloane, The On-Line Encyclopedia of Integer Sequences, A138158.

[17] V. Strehl, *Minimal transitive products of transpositions – the reconstruction of a proof of A. Hurwitz*, Sém. Lothar. Combin. **37** (1996), Art. S37c, 12 pp. (electronic).

Department of Mathematics, Bar-Ilan University, 52900 Ramat-Gan, Israel
E-mail address: `radin@math.biu.ac.il`

Department of Mathematics, Bar-Ilan University, 52900 Ramat-Gan, Israel
E-mail address: `yuvalr@math.biu.ac.il`